



TITLE:

# INVERSES OF A FAMILY OF BOUNDED LINEAR OPERATORS (Common Ground between Functional Analysis and Mathematical Theory of Information)

AUTHOR(S):

Saitoh, Saburou

---

CITATION:

Saitoh, Saburou. INVERSES OF A FAMILY OF BOUNDED LINEAR OPERATORS (Common Ground between Functional Analysis and Mathematical Theory of Information). 数理解析研究所講究録 2002, 1253: 152-163

ISSUE DATE:

2002-04

URL:

<http://hdl.handle.net/2433/41862>

RIGHT:

# INVERSES OF A FAMILY OF BOUNDED LINEAR OPERATORS

君美馬大・工 斎藤三郎 (Saburou Saitoh)

Department of Mathematics, Faculty of Engineering  
Gunma University, Kiryu 376-8515, Japan  
E-mail: ssaitoh@math.sci.gunma-u.ac.jp

## Abstract

We considered a generalization of the Pythagorean theorem with geometric meanings and from the generalization it seems that we were able to obtain a general and fundamental concept for the inversion of a family of bounded linear operators on a Hilbert space into various Hilbert spaces. After reviewing the applications to linear transforms in the framework of Hilbert spaces of the general theory of reproducing kernels, we shall state the results for the case of operator versions. In the last, we shall add the prototype example and meaning of the operator versions by figures, which show clearly a generalization of the Pythagorean theorem.

## 1. REPRODUCING KERNELS

We consider any positive matrix  $K(p, q)$  on  $E$ ; that is, for an abstract set  $E$  and for a complex-valued function  $K(p, q)$  on  $E \times E$ , it satisfies that for any finite points  $\{p_j\}$  of  $E$  and for any complex numbers  $\{C_j\}$ ,

$$\sum_j \sum_{j'} C_j \overline{C_{j'}} K(p_{j'}, p_j) \geq 0.$$

Then, by the fundamental theorem by Moore–Aronszajn, we have:

**Proposition 1.1**([1]) *For any positive matrix  $K(p, q)$  on  $E$ , there exists a uniquely determined functional Hilbert space  $H_K$  comprising functions  $\{f\}$  on  $E$  and admitting the reproducing kernel  $K(p, q)$  (RKHS  $H_K$ ) satisfying and characterized by*

$$K(\cdot, q) \in H_K \text{ for any } q \in E \tag{1.1}$$

and, for any  $q \in E$  and for any  $f \in H_K$

$$f(q) = (f(\cdot), K(\cdot, q))_{H_K}. \quad (1.2)$$

For some general properties for reproducing kernel Hilbert spaces and for various constructions of the RKHS  $H_K$  from a positive matrix  $K(p, q)$ , see the recent book [14] and its Chapter 2, Section 5, respectively.

## 2. CONNECTION WITH LINEAR TRANSFORMS

Let us connect linear transforms in the framework of Hilbert spaces with reproducing kernels ([7]).

For an abstract set  $E$  and for any Hilbert (possibly finite-dimensional) space  $H$ , we shall consider an  $H$ -valued function  $\mathbf{h}$  on  $E$

$$\mathbf{h} : E \longrightarrow H \quad (2.1)$$

and the linear transform for  $H$

$$f(p) = (\mathbf{f}, \mathbf{h}(p))_H \quad \text{for } \mathbf{f} \in H \quad (2.2)$$

into a linear space comprising functions on  $E$ . For this linear transform (2.2), we form the positive matrix  $K(p, q)$  on  $E$  defined by

$$K(p, q) = (\mathbf{h}(q), \mathbf{h}(p))_H \quad \text{on } E \times E. \quad (2.3)$$

Then, we have the following fundamental results:

(I) For the RKHS  $H_K$  admitting the reproducing kernel  $K(p, q)$  defined by (2.3), the images  $\{f(p)\}$  by (2.2) for  $H$  are characterized as the members of the RKHS  $H_K$ .

(II) In general, we have the inequality in (2.2)

$$\|f\|_{H_K} \leq \|\mathbf{f}\|_H, \quad (2.4)$$

however, for any  $\mathbf{f} \in H_K$  there exists a uniquely determined  $\mathbf{f}^* \in H$  satisfying

$$f(p) = (\mathbf{f}^*, \mathbf{h}(p))_H \quad \text{on } E \quad (2.5)$$

and

$$\|f\|_{H_K} = \|\mathbf{f}^*\|_H. \quad (2.6)$$

In (2.4), the isometry holds if and only if  $\{\mathbf{h}(p); p \in E\}$  is complete in  $H$ .

(III) We can obtain the inversion formula for (2.2) in the form

$$f \longrightarrow \mathbf{f}^*, \quad (2.7)$$

by using the RKHS  $H_K$ .

However, this inversion formula will depend on, case by case, the realizations of the RKHS  $H_K$ .

(IV) Conversely, if we have an isometric mapping  $\tilde{L}$  from a RKHS  $H_K$  admitting a reproducing kernel  $K(p, q)$  on  $E$  onto a Hilbert space  $H$ , then the mapping is linear and its isometrical inversion  $\tilde{L}^{-1}$  is represented in the form (2.2). Here, the Hilbert space  $H$ -valued function  $h$  satisfying (2.1) and (2.2) is given by

$$h(p) = \tilde{L}K(\cdot, p) \text{ on } E \quad (2.8)$$

and, then  $\{h(p); p \in E\}$  is complete in  $H$ .

When (2.2) is isometrical, sometimes we can use the isometric mapping for a realization of the RKHS  $H_K$ , conversely—that is, if the inverse  $L^{-1}$  of the linear transform (2.2) is known, then we have  $\|f\|_{H_K} = \|L^{-1}f\|_H$ .

We shall state some general applications of the results (I)~(IV) to several wide subjects and their basic references:

(1) Linear transforms ([7],[11]).

The fact that the image spaces of linear transforms in the framework of Hilbert spaces are characterized as reproducing kernel Hilbert spaces defined by (2.3) is the most important one in the general theory of reproducing kernels. Therefore, the fact will mean that the theory of reproducing kernels is fundamental and a general concept in mathematics. To look for the characterization of the image space is a starting point when we consider the linear equation (2.2). (II) gives a generalization of the Pythagorean theorem (see also [6]) and means that in the general linear mapping (2.2) there exists essentially an isometric identity between the input and the output. (III) gives a generalized (natural) inverse (solution) of the linear mapping (equation) (2.2). (IV) gives a general method determining and constructing the linear system from an isometric relation between outputs and inputs by using the reproducing kernel in the output space.

(2) Integral transforms among smooth functions ([18]).

We considered linear mappings in the framework of Hilbert spaces, however, we can consider linear mappings in the framework of Hilbert spaces comprising smooth functions, similarly. Conversely, reproducing kernel Hilbert spaces are considered as the images of some Hilbert spaces by considering some decomposed representations (2.3) of the reproducing kernels. Such decomposition is, in general, possible. This idea is important in [18] and also in the following items (6) and (7).

(3) Nonharmonic integral transforms ([8]).

If the linear system vectors  $h(p)$  move in a small way (perturbation of the linear system) in the Hilbert space  $H$ , then we can not calculate the related positive matrix (2.3), however, we can discuss the inversion formula and an isometric identity of the linear mapping. The prototype result is the Paley-Wiener theorem on nonharmonic Fourier series.

(4) Various norm inequalities ([8],[12]).

Relations among positive matrices correspond to those of the associated reproducing kernel Hilbert spaces, by the minimum principle. So, we can derive various norm inequalities among reproducing kernel Hilbert spaces. We were able to derive many beautiful norm inequalities.

(5) Nonlinear transforms ([12],[15]).

In a very general nonlinear transform of a reproducing kernel Hilbert space, we can look for a natural reproducing kernel Hilbert space containing the image space and furthermore, we can derive a natural norm inequality in the nonlinear transform. How to catch nonlinearity in connection with linearity? It seems that the theory of reproducing kernels gives a fundamental and interesting answer for this question.

(6) Linear integral equations ([19]).

(7) Linear differential equations with variable coefficients ([19]).

In linear integro-differential equations with general variable coefficients, we can discuss the existence and construction of the solutions, if the solutions exist. This method is called a backward transformation method and by reducing the equations to Fredholm integral equations of the first type -(2.2)- and we can discuss the classical solutions, in very general linear equations.

(8) Approximation theory ([3],[2]).

Reproducing kernel Hilbert spaces are very nice function spaces, because the point evaluations are continuous. Then, the reproducing kernels are a fundamental tool in the related approximation theory.

(9) Representations of inverse functions ([13]).

For any mapping, we discussed the problem of representing its inverse in term of the direct mapping and we derived a unified method for this problem. As a simple example, we can represent the Taylor coefficients of the inverse of the Riemann mapping function on the unit disc on the complex plane in terms of the Riemann mapping functions. This fact was important in the representation of analytic function in terms of local data in ([22],[23]).

- (10) Various operators among Hilbert spaces ([16]).

Among various abstract Hilbert spaces, we can introduce various operators of sum, product, integral and derivative by using the linear mapping (2.2) or very general nonlinear transforms. The prototype operator is convolution and we discussed it from a wide and general viewpoint with concrete examples.

- (11) Sampling theorems ([14], Chapter 4, Section 2; [5]).

The Whittaker-Kotel'nikov-Shannon sampling theorem may be interpreted by (I) and (II), very well and we can discuss the truncation errors in the sampling theory. J. R. Higgins [5] established a fully general theory for [14].

- (12) Interpolation problems of Pick-Nevanlinna type ([8],[9]).

General and abstract theory of interpolation problems of Pick-Nevanlinna type may be discussed by using the general theory of reproducing kernels.

- (13) Analytic extension formulas and their applications ([20],[10]).

We were able to obtain various analytic extension formulas and their applications from various isometrical identities (II). For their applications to nonlinear partial differential equations, see the survey article by N. Hayashi [4].

In this survey article, we shall present also new results on

- (14) Inversions of a family of bounded linear operators on a Hilbert space into various Hilbert spaces,  
which are generalizations of [21] and [6].

### 3. OPERATOR VERSIONS

We shall give operator versions of the fundamental theory (I) ~ (IV) which may be expected to have many concrete applications. In particular, for full generalizations of the Pythagorean theorem with geometric meanings, see [6]. Some special versions were given in [21].

For an abstract set  $\Lambda$ , we shall consider an operator-valued function  $L_\lambda$  on  $\Lambda$ ,

$$\Lambda \longrightarrow L_\lambda \tag{3.1}$$

where  $L_\lambda$  are bounded linear operators from a Hilbert space  $H$  into various Hilbert spaces  $\mathbf{H}_\lambda$ ,

$$L_\lambda : H \longrightarrow \mathbf{H}_\lambda. \tag{3.2}$$

In particular, we are interested in the inversion formula

$$L_\lambda x \longrightarrow x, \quad x \in H. \tag{3.3}$$

Here, we consider  $\{L_\lambda x; \lambda \in \Lambda\}$  as informations obtained from  $x$  and we wish to determine  $x$  from the informations. However, the informations  $L_\lambda x$  belong to various Hilbert spaces  $\mathbf{H}_\lambda$ , and so, in order to unify the informations in a sense, we shall take fixed elements  $\mathbf{b}_{\lambda,\omega} \in \mathbf{H}_\lambda$  and consider the linear mapping from  $H$

$$\begin{aligned} X_{\mathbf{b}}(\lambda, \omega) &= (L_\lambda x, \mathbf{b}_{\lambda,\omega})_{\mathbf{H}_\lambda} \\ &= (x, L_\lambda^* \mathbf{b}_{\lambda,\omega})_H, \quad x \in H \end{aligned} \quad (3.4)$$

into a linear space comprising functions on  $\Lambda \times \Omega$ . For the informations  $L_\lambda x$ , we shall consider  $X_{\mathbf{b}}(\lambda, \omega)$  as observations (measurements, in fact) for  $x$  depending on  $\lambda$  and  $\omega$ . For this linear transform (3.4), we form the positive matrix  $K_{\mathbf{b}}(\lambda, \omega; \lambda', \omega')$  on  $\Lambda \times \Omega$  defined by

$$\begin{aligned} K_{\mathbf{b}}(\lambda, \omega; \lambda', \omega') &= (L_{\lambda'}^* \mathbf{b}_{\lambda',\omega'}, L_\lambda^* \mathbf{b}_{\lambda,\omega})_H \\ &= (L_\lambda L_{\lambda'}^* \mathbf{b}_{\lambda',\omega'}, \mathbf{b}_{\lambda,\omega})_{\mathbf{H}_\lambda} \quad \text{on } \Lambda \times \Omega. \end{aligned} \quad (3.5)$$

Then, as in (I)  $\sim$  (IV), we have the following fundamental results:

(I') For the RKHS  $H_{K_{\mathbf{b}}}$  admitting the reproducing kernel  $K_{\mathbf{b}}(\lambda, \omega; \lambda', \omega')$  defined by (3.5), the images  $\{X_{\mathbf{b}}(\lambda, \omega)\}$  by (3.4) for  $H$  are characterized as the members of the RKHS  $H_{K_{\mathbf{b}}}$ .

(II') In general, we have the inequality in (3.4)

$$\|X_{\mathbf{b}}\|_{H_{K_{\mathbf{b}}}} \leq \|x\|_H, \quad (3.6)$$

however, for any  $X_{\mathbf{b}} \in H_{K_{\mathbf{b}}}$  there exists a uniquely determined  $x' \in H$  satisfying

$$X_{\mathbf{b}}(\lambda, \omega) = (x', L_\lambda^* \mathbf{b}_{\lambda,\omega})_H \quad \text{on } \Lambda \times \Omega \quad (3.7)$$

and

$$\|X_{\mathbf{b}}\|_{H_{K_{\mathbf{b}}}} = \|x'\|_H. \quad (3.8)$$

In (3.6), the isometry holds if and only if  $\{L_\lambda^* \mathbf{b}_{\lambda,\omega}; (\lambda, \omega) \in \Lambda \times \Omega\}$  is complete in  $H$ .

(III') We can obtain the inversion formula for (3.4) and so, for the mapping (3.3) as in (III), in the form

$$L_\lambda x \longrightarrow (L_\lambda x, \mathbf{b}_{\lambda,\omega})_{\mathbf{H}_\lambda} = X_{\mathbf{b}}(\lambda, \omega) \longrightarrow x', \quad (3.9)$$

by using the RKHS  $H_{K_{\mathbf{b}}}$ .

(IV') Conversely, if we have an isometric mapping  $\tilde{L}$  from a RKHS  $H_{K_{\mathbf{b}}}$  admitting a reproducing kernel  $K_{\mathbf{b}}(\lambda, \omega; \lambda', \omega')$  on  $\Lambda \times \Omega$  in the form (3.5) using

bounded linear operators  $L_\lambda$  and fixed vectors  $\mathbf{b}_{\lambda,\omega}$  onto a Hilbert space  $H$ , then the mapping  $\tilde{L}$  is linear and the isometric inversion  $\tilde{L}^{-1}$  is represented in the form (3.4) by using

$$L_\lambda^* \mathbf{b}_{\lambda,\omega} = \tilde{L} K_{\mathbf{b}}(\cdot, \cdot; \lambda, \omega) \text{ on } \Lambda \times \Omega. \quad (3.10)$$

Further, then  $\{L_\lambda^* \mathbf{b}_{\lambda,\omega}; (\lambda, \omega) \in \Lambda \times \Omega\}$  is complete in  $H$ .

## Acknowledgments

The author wishes to express my thanks Professor J. Kawabe for his kind invitation to the research meeting.

## References

- [1] N. Aronszajn. Theory of reproducing kernels. *Trans. Amer. Math. Soc.*, **68**(1950), 337–404.
- [2] D.-W. Byun and S. Saitoh. Approximation by the solutions of the heat equation. *J. Approximation Theory*, **78** (1994), 226–238.
- [3] D.-W. Byun and S. Saitoh. Best approximation in reproducing kernel Hilbert spaces. *Proc. of the 2th International Colloquium on Numerical Analysis, VSP-Holland*, (1994), 55–61.
- [4] N. Hayashi. Analytic function spaces and their applications to nonlinear evolution equations. *Analytic Extension Formulas and their Applications*, (2001), Kluwer Academic Publishers, 59–86.
- [5] J. R. Higgins. A sampling principle associated with Saitoh's fundamental theory of linear transformations. *Analytic Extension Formulas and their Applications*, (2001), Kluwer Academic Publishers, 73–86.
- [6] Th. M. Rassias and S. Saitoh. The Pythagorean theorem and linear mappings. *PanAmerican Math. J.* (to appear).
- [7] S. Saitoh. Hilbert spaces induced by Hilbert space valued functions. *Proc. Amer. Math. Soc.*, **89** (1983), 74–78.
- [8] S. Saitoh. Theory of Reproducing Kernels and its Applications. *Pitman Research Notes in Mathematics Series*, **189**(1988), Longman Scientific & Technical, UK.
- [9] S. Saitoh. Interpolation problems of Pick-Nevanlinna type. *Pitman Research Notes in Mathematics Series*, **212**(1989), 253–262.




- [10] S. Saitoh. Representations of the norms in Bergman-Selberg spaces on strips and half planes. *Complex Variables*, **19** (1992), 231–241.
- [11] S. Saitoh. One approach to some general integral transforms and its applications. *Integral Transforms and Special Functions*, **3** (1995), 49–84.
- [12] S. Saitoh. Natural norm inequalities in nonlinear transforms. *General Inequalities* **7**, (1997), 39–52. Birkhäuser Verlag, Basel, Boston.
- [13] S. Saitoh. Representations of inverse functions. *Proc. Amer. Math. Soc.*, **125** (1997), 3633–3639.
- [14] S. Saitoh. Integral Transforms, Reproducing Kernels and their Applications. *Pitman Research Notes in Mathematics Series*, **369** (1997). Addison Wesley Longman, UK.
- [15] S. Saitoh. Nonlinear transforms and analyticity of functions. *Nonlinear Mathematical Analysis and Applications*, (1998), 223–234. Hadronic Press, Palm Harbor.
- [16] S. Saitoh. Various operators in Hilbert space induced by transforms. *International J. of Applied Math.*, **1** (1999), 111–126.
- [17] S. Saitoh. Applications of the general theory of reproducing kernels. *Reproducing Kernels and their Applications*, (1999), Kluwer Academic Publishers, 165–188.
- [18] S. Saitoh and M. Yamamoto. Integral transforms involving smooth functions. *Reproducing Kernels and their Applications*, (1999), Kluwer Academic Publishers, 149–164.
- [19] S. Saitoh. Linear integro-differential equations and the theory of reproducing kernels. *Volterra Equations and Applications*. C. Corduneanu and I.W. Sandberg (eds), Gordon and Breach Science Publishers (2000), Amsterdam.
- [20] S. Saitoh. Analytic extension formulas, integral transforms and reproducing kernels. *Analytic Extension Formulas and their Applications*, (2001), Kluwer Academic Publishers, 207–232.
- [21] S. Saitoh. Applications of the reproducing kernel theory to inverse problems. *Comm. Korean Math. Soc.*, **16** (2001), 371–383.
- [22] S. Saitoh and M. Mori. Representations of analytic functions in terms of local values by means of the Riemann mapping function. *Complex Variables*, (to appear).
- [23] S. Saitoh. Principle of telethroscope. *Proceedings of the Graz Workshop on "Functional-analytic and complex methods, their interaction and applications to Partial Differential Equations"* (to appear).


The Pythagorean theorem:

$$\|x\|^2 = x_1^2 + x_2^2 ;$$

$$x_j = (x, e_j)$$



Generalizations of the Pythagorean theorem in the three types:




$$\begin{aligned} \|x\|^2 &= f_1(x_1, x_2) \\ &= f_2(y_1, y_2) \\ &= f_3(z_1, z_2) \end{aligned}$$

are valid for some functions  $f_1, f_2, f_3$  respectively:

$$x_j = (x, b_j), \quad x = y_1 b_1 + y_2 b_2,$$

$$z_j = \|x - (x, b_j) b_j\|.$$

However,  
 for  $n \geq 3$ ,  
 $\|x\|^2$   
 $= f(z_1, z_2, z_3)$   
 is impossible,  
 for any  
 function  $f$ .





However, we can establish an  
 isometric identity between

$$x \longleftrightarrow \left\{ x - (x, b_j) b_j \right\}_{j=1}^n ;$$

that is, when we consider  $\{z_j\}_{j=1}^n$  as  
 vectors.

Our operator versions give an  
 isometric identity in the linear  
 mapping and its inversion formula.





$$x = P_1 x + x_1$$

$$x_1 = P_2 x_1 + x_2$$

$$x_2 = P_1 x_2 + x_3$$

⋮

Even  $\lambda = 2$ , we, in general, need infinitely many procedures, if we wish to determine  $x$  by using two projections, step by step.